

COMUNICAZIONI

1) ANTIMICROBIAL SUSCEPTIBILITY OF SALMONELLA SPP. STRAINS ISOLATED FROM LAYER HENS IN CAMPANIA REGION FROM 2000 TO 2003

Ludovico Dipineto, Claudia Scarpetta, Mariarosaria Calabria, Mariangela Sensale, Antonio Baiano, Lucia Francesca Menna, Alessandro Fioretti

Dipartimento di Patologia e Sanità Animale, Università di Napoli Federico II, Italy

Corresponding author: Prof. Alessandro Fioretti, Dipartimento di Patologia e Sanità Animale. Facoltà di Medicina Veterinaria, Università di Napoli Federico II. Via Delpino, 1, 80137 Napoli (NA), Italy - Tel. +39 081451802 - Fax: +39 0815091993 - Email: fioretti@unina.it

Abstract The aim of this study was to determine the antimicrobial resistance in 60 Salmonella strains (*S. enteritidis*, *S. typhimurium*, *S. gallinarum*) isolated from layer hens in Campania region from 2000 to 2003. *S. gallinarum* showed resistance against ciprofloxacin and enrofloxacin, in contrast, *S. enteritidis* and *S. typhimurium* were fully susceptible. In all of isolates high levels of resistance were observed for neomycin, gentamycin and oxytetracycline. Also, one significant observation was that all of the isolates showed full susceptibility to Sulphamethoxazole/Trimethoprim.

These results suggest importance to restrict the use of antibiotics in layers hens flocks in order to reduce the selection and spread of multiresistant strains

Key words: Antimicrobial resistance, Salmonella strains

Sensibilità agli Antibiotici di Salmonella spp. Isolati da Galline Ovaiole in Campania nel Triennio 2000/2003

Riassunto Scopo del presente lavoro è stato quello di testare la sensibilità antibiotica di 60 ceppi di Salmonella (*S. enteritidis*, *S. typhimurium*, *S. gallinarum*) isolati da galline ovaiole nel periodo compreso tra il 2000 e il 2003. *S. gallinarum* mostrava resistenza nei confronti di ciprofloxacina ed enrofloxacina (rispettivamente 15% e 23%), al contrario di *S. enteritidis* e *S. typhimurium* che manifestavano una completa sensibilità. Tutti i sierotipi valutati presentavano alte percentuali di resistenza nei confronti di neomicina, gentamicina e ossitetraciclina. Nei confronti dei sulfamidici i ceppi testati presentavano resistenza nulla. Alla luce di quanto esposto si consiglia un uso più moderato e mirato degli antibiotici negli allevamenti in modo da ridurre la selezione e diffusione di ceppi multiresistenti.

Parole chiave: resistenza antimicrobica, Salmonelle

Introduction

Antimicrobial resistance is the capacity of bacteria to survive exposure to a defined concentration of an antimicrobial substance. It is the natural response of bacterium to defend itself against the effects of an antibiotic. The development of antimicrobial resistance is an ecological phenomenon. Any antibiotic use, whether in humans, animals or plants/environment may lead to resistance (OIE, 2003).

The extensive use of antibiotics, not only in human and veterinary medicine, but also in livestock production for disease prevention or as growth-promoting feed additives, has led to a serious increase in, and spread of, multiple antibiotic-resistant bacteria (Cruchaga et al., 2001). All this caused considerable problems to approach prophylactics and therapeutics plans versus various bacterial pathologies.

Generally, the increased application of antimicrobials in veterinary and human medicine has been implicated as a contributing factor in the emergence of antimicrobial-resistant pathogens and the evolution of multiple drug-resistant strains.

Salmonellosis has a particular role in avian medicine whether host-specific serotypes (*S. gallinarum*, *S. pullorum*) or non-host specific serotypes (*S. enteritidis*, *S. typhimurium*) implicated in foodborne zoonoses.

Within the routine control programmes carried out in the poultry farms organized from the avian pathology section of the Dipartimento di Patologia e Sanità Animale; the Salmonella strains isolated showed an increase of antibiotic resistance pattern.

The aim of this study was to determine the antimicrobial resistance in Salmonella strains (*S. enteritidis*, *S. typhimurium*, *S. gallinarum*) isolated from layer hens in Campania region from 2000 to 2003.

Materials and methods

Sample collection

From January 2000 to November 2003 a total of 60 *Salmonella* strains were isolated from layer hens flocks, respectively belonged to *S.gallinarum*, *S.enteritidis*, *S.typhimurium*. The strains collected were 20 for each serotype.

Isolation and identification procedure

The *Salmonella* isolation procedures were carried out following the WHO standard methods (WHO, 1994). All the strains were serotyped at National Reference Centre for *Salmonella* (Istituto Zooprofilattico Sperimentale delle Venezie, Padova - Italy).

Antimicrobial susceptibility tests

Antimicrobial susceptibility profiles of the isolates were determined by the disk diffusion method according to the NCCLS guidelines (National Committee for Clinical Laboratory Standards, 1997). The antimicrobial agents (Oxoid) tested and corresponding concentration were as follows: Ciprofloxacin 5µg (C), Enrofloxacin 5µg (E), Flumequine 30µg (F), Nalidixic acid 30µg (NA), Apramycin 15µg (AP), Amoxicillin 10µg (A), Neomycin 30µg (N), Gentamicin 10µg (G), Oxytetracycline 30µg (O), Sulphamethoxazole/Trimethoprim 25µg (S/T).

The diameters of the inhibition zone for the interpretation of resistance and susceptibility were those recommended by the NCCLS (National Committee for Clinical Laboratory, 2002). Results were scored as susceptible, moderately susceptible or resistant according to NCCLS criteria (2002). The *Escherichia coli* ATCC 25 922 was used as reference strain.

Results

As seen from Table 1, resistance of *S. gallinarum* was significantly higher than other two serotypes examined. In particular, *S. gallinarum* showed resistance against two fluoroquinolone (ciprofloxacin and enrofloxacin, respectively 15% and 23%), in contrast, *S. enteritidis* and *S. typhimurium* were fully susceptible. In all of isolates high levels of resistance were observed for neomycin, gentamycin and oxytetracycline. It was also found that *S. enteritidis* and *S. gallinarum* were resistant to apramycin (33,3% and 38,5% respectively) and *S. gallinarum* was resistant to amoxicillin too (23,1%). In conclusion, one significant observation was that all of isolates showed full susceptibility to Sulphamethoxazole/Trimethoprim.

Table 1. Antimicrobial resistance of 60 *Salmonella* spp. isolates.

Serotype	n°	Antimicrobial resistance (%)									
		C	E	NA	AP	A	N	G	F	S/T	O
<i>S. enteritidis</i>	20	0	0	6	33,3	0	55,5	38,8	5,5	0	33,3
<i>S. typhimurium</i>	20	0	0	13	0	0	12,5	25	12,5	0	50
<i>S. gallinarum</i>	20	15	23	15	38,5	23,1	53,8	55,3	23,1	0	46,1

Discussion

An increase in the incidence of antibiotic resistance in *Salmonella* isolated from humans and animals related to exhaustive application of antibiotics in both groups has been documented worldwide (Chruchaga et al., 2001). Recently, Lee et al. (Lee et al., 2003) reported, in an antimicrobial susceptibility test against 258 isolates of *S.gallinarum*, a reduced susceptibility to ampicillin (13.0%), gentamicin (43.4%), kanamycin (69.6%), enrofloxacin (6.5%), ciprofloxacin (10.9%), norfloxacin (52.5%) and ofloxacin (82.6%). A study on antimicrobial-resistant *Salmonella enterica* serovars isolated from chickens in Spain showed high percentage of resistance to chloramphenicol (44.6%), ampicillin (34.8%) and tetracycline (33.9%) (Hernandez et al., 2002). Jones et al. (2002) reported *S.typhimurium* strains resistant to ampicillin, sulphonamides, streptomycin, chloramphenicol and tetracyclines as well as *S.typhimurium* isolated from poultry resistant to nalidixic acid (Jones et al., 2002).

Fluoroquinolones resistance was rarely found among *Salmonella* species until Heisig reported *S.typhimurium* serovar *Copenhagen* from cattle was highly resistant to ciprofloxacin (Heisig, 1993).

Conclusions

An important finding is the antimicrobial resistance observed in *S.gallinarum* against fluoroquinolones (ciprofloxacin and enrofloxacin). High percentage of resistance observed in *S.typhimurium* and *S.enteritidis* for neomycin, gentamycin and oxytetracycline demonstrate improper use of these antibiotics in the control of avian salmonellosis, particularly in metaphylactic sense.

These results confirm importance to restrict the use of antibiotics in layers hens flocks in order to reduce the selection and spread of multiresistant strains and underlines the need for integrated surveillance systems of antibiotic resistance that consider isolates not only from human disease but also from the animal reservoirs and the food vehicles.

Acknowledgements

We would like to thank Dr. Antonia Ricci, National Reference Centre for Salmonella - Istituto Zooprofilattico Sperimentale delle Venezie, Padova, Italia, for the serotyping of our isolates and Mrs. Fortuna Pisa for her collaboration.

References

1. Cruchaga, S., Echeita, A., Aladuena, A., Garcia-Pena, J., Frias, N., Usera, M. A., 2001. Antimicrobial resistance in salmonellae from humans, food and animals in Spain in 1998. *J. Antimicrob. Chemother.* 47(3):315-21.
2. Heisig, P., 1993. High-level fluoroquinolone resistance in a *Salmonella typhimurium* isolate due to alterations in both *gyrA* and *gyrB* genes. *J. Antimicrob. Chemother.* 32(3):367-77.
3. Hernandez, T., Rodriguez-Alvarez, C., Arevalo, M. P., Torres, A., Sierra, A., Arias, A., 2002. Antimicrobial-resistant *Salmonella enterica* serovars isolated from chickens in Spain. *J. Chemother.* 14(4):346-50.
4. Jones, Y. E., Chappell, S., McLaren, I. M., Davies, R. H., Wray, C., 2002. Antimicrobial resistance in *Salmonella* isolated from animals and their environment in England and Wales from 1988 to 1999. *Vet. Rec.* 25;150(21):649-54.
5. Lee, Y. J., Kim, K. S., Kwon, Y. K., Tak, R. B., 2003. Biochemical characteristics and antimicrobials susceptibility of *Salmonella gallinarum* isolated in Korea. *J. Vet. Sci.* 4(2):161-6.
6. NCCLS (National Committee for Clinical Laboratory Standards), 2002. Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals, second edition: approved standard M31-A2. Wayne, PA, USA.
7. OIE (Office International des Épizooties), 2003. OIE International Standards on Antimicrobial Resistance. Paris, France.
8. WHO, 1994. Guidelines on detection and monitoring of *Salmonella* infected poultry flocks with particular reference to *Salmonella enteritidis*. In: Wray, C., Davies, R. H., (eds.), WHO, Veterinary Public Health Unit.

2) PREVALENCE OF *CAMPYLOBACTER JEJUNI* IN POULTRY BREEDER FLOCKS

Lucia Francesca Menna¹, Gianluca Matteoli¹, Marzia Fontanella¹, Alessandra Cuomo¹, Antonio De Paola², Tiziana Pepe³, Isolina Di Marco³, Ludovico Dipineto¹

¹ Dipartimento di Patologia e Sanità Animale, Università di Napoli Federico II, Italy,

² Arena Holding Spa, Loc. Monteverde, Bojano (CB), Italy,

³ Dipartimento di Scienze Zootecniche ed Ispezione degli Alimenti, Università di Napoli Federico II, Italy

Corresponding author: Prof. Lucia Francesca Menna, Dipartimento di Patologia e Sanità Animale. Facoltà di Medicina Veterinaria, Università di Napoli Federico II. Via Delpino, 1, 80137 Napoli (NA), Italy - Tel. +39 081451802 - Fax: +39 0815091993 - Email: fioretti@unina.it

Abstract The aim of this work is to present the preliminary results of a study about breeders carrying out in order to demonstrate the supposed correlation between *Campylobacter* isolated from breeders and the ones isolated from broilers, supporting the theory of the vertical transmission of the germ. It was examined three different breeder flocks of Bojano in Molise region.

A total of 360 cloacal swabs and 80 environmental swabs was collected. Of the 3 flocks studied, 6,9% tested were positive for *Campylobacter spp.* The most-prevalent isolated species is *C. jejuni* (8,2%). Only 3 of the 360 cloacal swabs samples examined were associated with *C. coli*. The environmental swabs resulted negative. This results confirms again that poultry is a reservoir of this germ.

Indagine sulla Prevalenza di *Campylobacter jejuni* in Gruppi di Riproduttori Avicoli

Riassunto Il presente lavoro si propone di illustrare i risultati preliminari, sui riproduttori, di uno studio più ampio volto a stabilire, mediante studi di biologia molecolare, una eventuale correlazione tra i *Campylobacter* isolati dai riproduttori e quelli isolati dai broiler in modo da definire l'eventuale trasmissione verticale del germe. Sono stati esaminati 3 gruppi di riproduttori ubicati a Bajano in Molise. Dei tre gruppi valutati, il 6,9% risultava positivo a *Campylobacter spp.* *C. jejuni* era la principale specie isolata (8,2%). Solo 3 dei 360 tamponi cloacali esaminati era associata a *C. coli*. I tamponi ambientali risultavano negativi. Tali risultati confermano ancora una volta il ruolo del pollame come reservoir di questo microorganismo.